

Updated Criteria for Population-Based Stroke and Transient Ischemic Attack Incidence Studies for the 21st Century

Valery Feigin, MD, PhD; Bo Norrving, MD; Cathie L.M. Sudlow, FRSE;
Ralph L. Sacco, MD

Stroke is the second leading cause of death and disability worldwide and the burden of stroke over the last 3 decades has increased significantly.¹ There have also been noticeable changes in stroke epidemiology, including diverging trends in stroke incidence and mortality rates in high-income and low-to middle-income countries,² disproportionately increasing burden of stroke in developing countries,³ and increased proportion of disability and deaths because of stroke compared with all diseases.⁴ Moreover, although most countries have no accurate data on stroke incidence and outcomes,⁵ in countries where such studies have been conducted we know that there are large within and between country geographical differences in stroke burden,^{1,3} large ethnic/racial disparities in stroke,^{6–11} and noticeable secular changes in the natural history of stroke and risk factors.^{12,13}

All these changes, differences, trends, and gaps in knowledge necessitate further, more advanced epidemiological studies of stroke in various populations. Accurate and comparable population-based data on stroke incidence and outcomes and their trends can provide reliable data for evidence-based stroke care planning, healthcare resource allocation, and priority setting. Current criteria for population-based stroke incidence studies include a standard clinical World Health Organization (WHO) definition of stroke and transient ischemic attack (TIA)¹⁴ and maximally complete case ascertainment of both fatal and nonfatal, hospitalized and nonhospitalized new stroke events using multiple overlapping sources of information.^{15–20}

More recently, the definitions of TIA and stroke have been updated to incorporate both clinical and tissue criteria by the American Heart Association and American Stroke Association,²¹ and a similar definition also appears in the upcoming *International Classification of Diseases, Eleventh Revision (ICD-11)*.²² The new definitions of stroke and TIA require a reconsideration of the criteria for an ideal stroke incidence study. Moreover, there are more countries using/adapting alternative methods for population-based stroke incidence

and outcome estimates, including national health record-linkage systems and various stroke registries. The issue becomes even more imperative given ongoing advances in acute stroke management and rehabilitation that require additional accurate epidemiological and healthcare information from stroke incidence studies. The aims of this article are to extend and refine the latest criteria for an ideal stroke incidence study^{17,18} taking into consideration the new stroke and TIA definitions and latest developments in stroke care and systems with the goal of suggesting updated criteria for standard definitions, methods, and data presentation (Table 1).

Updated Standard Definitions of TIA and Stroke

The addition of the WHO clinical definition of TIA (any acute focal cerebrovascular event with symptoms lasting <24 hours)¹⁴ to the core criteria would allow evaluation of the burden of 2 major acute cerebrovascular disorders—stroke and TIA—separately and combined. A wide use of thrombolytic therapy within the first hours after acute ischemic cerebrovascular event occurs when the time-based clinical differentiation of stroke and TIA is not possible means that new TIA events could and should be registered along with new stroke events. From a public health perspective, it is important to know the incidence of all acute cerebrovascular events, including TIA, as they all require urgent diagnosis and management with associated healthcare resource utilization. This means that a search for new TIA events must be as complete and thorough as it is for new stroke events, although it will require additional resources and may be a challenging task in resource-poor countries. Additionally, adjudication of potential TIA events in a community setting is particularly challenging because of TIA mimics and hospitalization bias.^{23,24} However, there are already a few population-based studies in which both stroke and TIA were ascertained and registered at the same time in the same study population,^{25–30} thus providing support for the

Received May 10, 2018; final revision received June 6, 2018; accepted July 13, 2018.

From the National Institute for Stroke and Applied Neurosciences, School of Public Health and Psychosocial Studies, Faculty of Health and Environmental Sciences, Auckland University of Technology, New Zealand (V.F.); Department of Clinical Sciences, Department of Neurology, Skane University Hospital, Lund University, Sweden (B.N.); Centre for Medical Informatics, Usher Institute, University of Edinburgh, UK Biobank, United Kingdom (C.L.M.S.); and Department of Neurology, McKnight Brain Institute, Clinical & Translational Science Institute, Miller School of Medicine, University of Miami, FL (R.L.S.).

Guest Editor for this article was Emmanuel Touzé, PhD.

Correspondence to Valery Feigin, MD, PhD, National Institute for Stroke and Applied Neurosciences, School of Public Health and Psychosocial Studies, Faculty of Health and Environmental Sciences, AUT University, 90 Akoranga Dr, Northcote, Auckland 0627, New Zealand. Email valery.feigin@aut.ac.nz (Stroke. 2018;49:00-00. DOI: 10.1161/STROKEAHA.118.022161.)

© 2018 American Heart Association, Inc.

Stroke is available at <https://www.ahajournals.org/journal/str>

DOI: 10.1161/STROKEAHA.118.022161

Table 1. Criteria for Basic and Advanced Population-Based Stroke and TIA Incidence Studies

Domains	Core Criteria for Basic Studies	Supplementary Criteria for Advanced Studies
Standard definitions	WHO clinical definition of stroke	AHA/ASA/ICD-11 clinical and tissue definition of stroke*
	WHO clinical definition of TIA*	TIA without evidence of brain infarct*
	At least 80% CT/MRI verification of the diagnosis of ischemic stroke, intracerebral hemorrhage, and subarachnoid hemorrhage within the first week of stroke onset	At least 80% DW-MRI for tissue-based distinction of TIA and stroke for events resolving within first 24 h of the onset*
	First-ever-in-a-lifetime stroke or TIA	Classification of ischemic stroke, intracerebral hemorrhage, and subarachnoid hemorrhage into etiologic and anatomic groups*
		Stroke severity by NIHSS or similar*
		Recurrent stroke and TIA*
Standard methods	Complete, population-based case ascertainment, based on multiple overlapping sources of information (hospitals, outpatient clinics, general practitioners, death certificates)	Ascertainment of patients with TIA, recurrent strokes, and those referred for brain, carotid, or cerebral vascular imaging
	Prospective study design	Hot pursuit of cases
	Large, well-defined, and stable population, allowing at least 100 000 person-years of observation	Direct assessment of underascertainment by regular checking of general practitioners' databases and hospital admissions for acute vascular problems and cerebrovascular imaging studies and interventions
	Reliable method for estimating denominator (census data ≤ 5 -y-old)	Capture-recapture estimates of the completeness of case ascertainment*
	Follow-up of patients' vital status for at least 1 mo	Follow-up of stroke and TIA patients' functional status for at least 3 mo*
		Collecting data on stroke risk factors and quality of acute care*
Standard data presentation	Complete calendar years of data; ≤ 5 y of data averaged together	Data on old (WHO) and new (AHA/ASA/ICD-11) stroke and TIA estimates presented separately*
	Men and women presented separately	Data also presented by major ethnic/racial groups*
	Mid-decade 10- or 5-y age bands (eg, 55 to 64 y) used in publications, including oldest age group (≥ 85 y)	
	95% CI around rates	
	STROND guidelines for reporting*	

AHA/ASA indicates American Heart Association/American Stroke Association; CT, computed tomography; DW-MRI, MRI with diffusion-weighted sequences; ICD, International Classification of Diseases; MRI, magnetic resonance imaging; NIHSS, National Institutes of Health Stroke Scale; STROND, Standards of Reporting of Neurological Disorders; TIA, transient ischemic attack; and WHO, World Health Organization.

*New criteria.

Adapted from Feigin et al¹⁷ with permission. Copyright © 2004, American Heart Association, Inc. Adapted from Feigin et al¹⁸ with permission. Copyright © 2004, Elsevier.

feasibility of such joint stroke and TIA incidence studies in different populations.

Although it is crucial to keep using the previous WHO clinical definition of stroke (rapidly developing clinical signs of focal [or global] disturbance of cerebral function, lasting >24 hours or leading to death, with no apparent cause other than that of vascular origin)¹⁴ and TIA (a sudden onset of a focal neurological symptom and sign lasting <24 hours and caused by reversible cerebral ischemia)¹⁴ in all epidemiological studies of stroke incidence to allow comparisons between different locations and over time (geographical and trends analysis), it is also important to include/add the new tissue-based definition of stroke and TIA in the latest epidemiological studies. Routine use of modern neuroimaging, vascular/cardiac, and other laboratory investigations has become

generally available in many hospitals/centers, thus allowing more precise etiologic and anatomic classifications of TIA, ischemic stroke, intracerebral hemorrhage, and subarachnoid hemorrhage.

In the new American Heart Association/American Stroke Association definition, ischemic stroke is defined on the basis of clinical and tissue criteria as brain, spinal cord, or retinal cell death attributable to ischemia, based on neuropathological, neuroimaging, and clinical evidence of permanent injury.³¹ Effectively, it includes formerly defined TIAs with neuroimaging evidence of ischemic brain lesion in the clinically relevant area of the brain and does not depend on the duration of neurological symptoms associated with the cerebrovascular ischemic event. TIA is now defined as transient episode of neurological dysfunction caused by focal brain,

spinal cord, or retinal ischemia, without acute infarction,²¹ thereby eliminating the time-based components noted in the classical WHO definition.¹⁴ Moreover, the new American Heart Association/American Stroke Association definition for stroke included silent or subclinical tissue-based strokes that can only be detected with more sensitive brain imaging. Most population-based incidence studies would not be able to adequately enumerate silent strokes because of the need for advanced imaging and the inability to accurately determine the date of onset of such events.

In the upcoming *ICD-11* at WHO (<https://icd.who.int/dev11/l-m/en>), TIA is defined as transient episode of focal neurological dysfunction caused by focal brain ischemia without acute infarction in the clinically relevant area of the brain. Symptoms should resolve completely within 24 hours. Cerebral ischemic stroke is defined as acute focal neurological dysfunction caused by focal infarction at single or multiple sites of the brain. Evidence of acute infarction may come either from (1) symptom duration lasting >24 hours or (2) neuroimaging or other technique in the clinically relevant area of the brain. The term does not include infarction of the retina. Although the *ICD-11* definitions of stroke and TIA do not include monocular blindness with transient ischemia of the retina or retinal infarct, to be consistent with previous epidemiological studies of stroke we recommend to register these retinal ischemic events. The WHO *ICD-11* definition of stroke requires the presence of acute neurological dysfunction and encompasses the entities cerebral ischemic stroke, intracerebral hemorrhage, subarachnoid hemorrhage, and stroke not known to be ischemic or hemorrhagic. In the WHO *ICD-11* definition, silent cerebral infarcts and microinfarcts are not included in the definition of stroke.

Both new definitions require more advanced neuroimaging or neuropathological confirmation of infarction. Computerized tomography is less sensitive to detection of small ischemic lesions than magnetic resonance imaging with diffusion-weighted sequences, which is more likely to detect an acute lesion in the clinically relevant area. Increasing use of magnetic resonance imaging with diffusion-weighted sequences will lead to transient cerebral events labeled as TIAs under the classical WHO definition being called cerebral ischemic strokes under the WHO *ICD-11* definition.

According to the American Heart Association/American Stroke Association stroke definitions, intracerebral hemorrhage is defined as rapidly developing clinical signs of neurological dysfunction attributable to a focal collection of blood within the brain parenchyma or ventricular system that is not caused by trauma; and subarachnoid hemorrhage as rapid signs of neurological dysfunction and headache because of bleeding into the subarachnoid space (the space between the arachnoid membrane and pia mater of the brain or spinal cord), which is not caused by trauma. These definitions of intracerebral hemorrhage and subarachnoid hemorrhage are similar to the *ICD-11* definitions.²²

To allow meaningful comparison of epidemiological studies by stroke etiologic and anatomic subtypes and over time it is important to use standard and validated classification systems. Ischemic stroke can be further classified into 5 etiologic groups (large-artery atherosclerosis, cardioembolism,

small-vessel occlusion, stroke of other determined etiology, and stroke of undetermined cause) according to the TOAST (Trial of Org 10172 in Acute Stroke Treatment) classification.³² If possible, stroke of undetermined cause can be further classified based on the extent of the diagnostic investigations into 3 groups: (1) ≥ 2 causes identified, (2) negative evaluation despite a complete investigation, and (3) incomplete diagnostic evaluation.³² The *ICD-11* classification of cerebral ischemic stroke builds on the TOAST classification. Some studies have also proposed classifying a proportion of the infarcts of undetermined cause (cryptogenic strokes) with adequate diagnostic investigations as embolic stroke of undetermined source among those with nonlacunar syndromes without significant proximal arterial atherosclerosis or a definite cardioembolic source.^{33,34} More recent etiologic classification schemes, such as the Clinical Classification System, have been developed with the aim of providing improvements in interobserver reliability and reducing the proportion of ischemic strokes attributed to undetermined cause.^{35,36} Clinical anatomic subgroups have also been used to classify ischemic strokes into 5 anatomic groups (total anterior circulation syndrome, partial anterior circulation syndrome, lacunar syndrome, posterior circulation syndrome, and uncertain circulation syndrome) according to the Oxfordshire Community Stroke Project clinical classification.³⁷ This classification system provides less information on presumed cause, but is less dependent on the availability of data from brain and vascular imaging, giving it the advantage that almost all strokes can be assigned to a subtype. Better comparability of studies by stroke severity are achieved through use of a well-validated and widely used scales, such as the National Institutes of Health Stroke Scale (NIHSS).³⁸ National Institutes of Health Stroke Scale score can be obtained either prospectively (if clearly present in medical files) or retrospectively based on clinical data obtained from medical files.³⁹ Such subtype classification and severity scoring systems are recommended to be used as supplementary criteria for an ideal stroke incidence and outcomes study.

Although categorization of different pathological types of stroke into various etiologic and anatomic groups might be very challenging in population-based epidemiological studies, especially in resource-poor countries, there have been several such studies accomplished in the past.^{40–49} For example, if possible, intracerebral hemorrhages should be classified into lobar, deep, and infratentorial hemorrhages; whereas subarachnoid hemorrhages should be classified into aneurysmal and nonaneurysmal hemorrhages. These classifications help in the management, and predicting prognosis and planning secondary preventive measures, justifying their use in population-based stroke incidence studies. Such classification systems may also require the use of an adjudication committee to reduce bias and help minimize misclassification.²³

Suggested Supplementary Data Collection

Suggested additional supplementary data collection also includes registration of recurrent TIA (because recurrent TIA has different management and prognostic significance); capture-recapture estimates to ascertain the completeness of stroke and TIA case ascertainment,^{50,51} follow-up of stroke and TIA patients' functional status for at least 3 months (there is




Figure. A basic population-based stroke and transient ischemic attack incidence study using door-to-door survey. Adapted from Feigin²⁰ with permission. Copyright © 2007, Elsevier.

evidence that 3-month disability, as measured by the modified Rankin Scale pre- and poststroke, is a strong independent predictor of long-term disability and mortality^{52,53}, and collecting data on risk factors among stroke and TIA patients to allow firmer conclusions to be drawn about the factors impacting the trends in incidence and outcomes as well as to reflect on the effect of trends in health policy on stroke and TIA outcomes.^{40,54} Complete, population-based case ascertainment, based on multiple overlapping sources of information (hospitals, outpatient clinics, general practitioners, death certificates) and high level of acute brain neuroimaging coverage, with an expert adjudication of the events is very important for the capture and diagnosis/classification of all new stroke and TIA cases²³ based on both old (clinical) and new stroke and TIA definitions. The new definitions of stroke and TIA require the use of magnetic resonance imaging with diffusion-weighted sequences for the precise distinction between TIA and ischemic stroke if the events are shorter than 24 hours, whereas for events lasting 24+ hours brain computed tomographic scanning is sufficient. Consistent with previous criteria for a population-based stroke incidence study,^{16–18} we recommend that when magnetic resonance imaging with diffusion-weighted sequences is done for cerebrovascular events lasting <24 hours it must be done in less than a week at least 80% of the time. Reporting of proportions examined and timing of the examination with each technique is important for comparative purposes.

Given the prognostic importance of stroke for the subsequent development of cognitive decline and dementia, as well as its effect on quality of life and other health outcomes, it is also recommended to include additional validated

measurements (eg, cognitive impairment, daily activities, quality of life, mood, socioeconomic status, and direct and indirect costs). Preferences should be given to the validated assessment tools that are easier (eg, over the phone) and faster to administer in stroke/TIA patients and their family caregivers. As quality and timing of acute stroke/TIA care are important determinants of their outcomes, it is also desirable to include some quality indicators of stroke care.⁵⁵ The choice and number of assessment tools to be used in a population-based stroke incidence study should be determined by the purpose of the study, expertise of study researchers, and available resources.

Updated Standard Methods of Case Ascertainment

The standard methods for conducting population-based stroke incidence studies are also applicable for determining stroke incidence using either the WHO STEPS (STEPwise Approach to Surveillance) approach to stroke surveillance,¹⁹ national registries (eg, record-linkage systems), or a special door-to-door survey²⁰ (Figure). These survey methods require use of instruments that allow clinical ascertainment of nonhospitalized and nonfatal stroke and TIA events among study participants and include a validated verbal-autopsy technique for fatal events (interviewing family members and relevant health professionals about the circumstances of a death after the event over the last couple of years). Such door-to-door surveys for studying stroke incidence have been successfully accomplished in Italy and China.^{56–58} Advantages of this approach include relatively low cost and simplicity (door-to-door survey is the most commonly used design of

stroke epidemiological studies in resource-poor settings). Disadvantages include the potential for underestimating incidence rates by missing very mild stroke cases, recall bias (although this can be minimized by restricting the period of recall to 3 years), and the need to study a large sample size.²⁰ It should be emphasized that the WHO STEPS stroke surveillance can only be called a population-based study if it includes all 3 steps of the surveillance (hospital register, death certificates, and nonhospitalized case ascertainment in the catchment area). The updated classifications of TIA and stroke may also require greater access to advanced brain

imaging to use classification systems that rely on tissue-based criteria for infarction and hemorrhage.

Updated Standard Data Presentation

To allow comparisons between different stroke incidence studies and analysis of temporal trends, it is important to conduct a study in a way that (1) allows accommodation of both core and supplementary criteria and (2) follows standard guidelines for reporting study results, such as the Standards of Reporting of Neurological Disorders checklist (Table 2).^{59,60} We suggest calling a study that meets both

Table 2. Standards for Reporting Population-Based Stroke and TIA Incidence Studies

Section/Topic	Recommendation
Title	Give the type of study design used using a widely recognized term.
Abstract	The abstract should give an accurate summary of how the study was conducted and the main findings.
Introduction	
Background	Details of the scientific rationale for the study should be reported.
Aims/objectives	State the specific aims and objectives of the study.
Methods	
Study design	Give a full description of the study design, and what criteria it meets to be called a population-based study.
Setting	Clearly defined (usually, but not always, on a geographic basis), and stable, with reliable information on in- and out-migration.
Source population	Description of how all eligible members of the population were identified and through what data sources (eg, hospitals, outpatient clinics, death certificates). Source of data used for the study (eg, administrative database, medical records). If administrative database used algorithms for data extraction should be described. Description of the rate of hospital admission. Details of healthcare system in the country (study region) where the study was conducted (eg, public versus private healthcare system). Description of how a person with stroke or TIA is referred in the country (study region) where the study was conducted. Description and characteristics of response rate/dropouts and exclusion rate if applicable.
Study participants	Definition of stroke and TIA is clearly identified and presented in sufficient detail. Details of the sampling method are described (are participants representative of the source population). Fully validated source of diagnosis or reference-standard criteria applied. Definition and justification of disease severity (preferably using a standardized scale). Description of how types/subtypes of stroke/TIA are distinguished (if relevant). Description of how completeness of case ascertainment was assessed. Description of whether completeness of case ascertainment was adequate.
Ethical approval	Details of ethics approval/informed consent/data governance should be reported.
Measurement	Give details of how incidence was determined (based on timing of data collection either prospectively or retrospectively). Definition and justification of timing of measurements. The data presented to some specified time period (usually whole years or person-time). Raw numbers are reported in sufficient detail to calculate the appropriate rates (eg, by age or sex).
Statistical methods	If rates have been standardized (eg, by age or sex), then the details of the standard population used should be given. If possible 2 standard populations should be used one with local relevance and the other to facilitate international comparisons. Description of any assumptions made in the calculations should be reported. An explanation of how missing data were addressed in the analyses.
Results	
Main findings	Consider a flow diagram that describes how participants were included in the study (useful to assess how a person with stroke/TIA is referred). Give appropriate rates with their associated 95% CIs. Report results of any sensitivity analyses.
Discussion	
Key findings	Summarized the key findings in relation to the study aims and objectives.
Interpretation	Interpret the results in the context of the evidence from other well-performed studies with similar designs and objectives. Reliability of the estimates (ie, based on the reporting of the statistical methodology, and study design, measurement of key information).
Limitations	Discuss potential limitations of the study. Include details of risk of bias (eg, selection bias), completeness of case ascertainment, and data quality (assessment of its probability, size, and potential importance).
Generalizability	Discuss the external validity of the study findings. Are the results consistent with reviews of descriptive epidemiological studies on the same topic that cover different settings?
Implications	Discuss implications of the research findings for practice and future research.

TIA indicates transient ischemic attack.

Adapted from Bennett et al⁵⁹ with permission. Copyright © 2015, Karger Publishers.

core and supplementary criteria and standard data presentation (Tables 1 and 2) an advanced population-based stroke incidence study. However, such stroke incidence studies are expensive and require special expertise for their design and execution, which is particularly challenging in resource-poor countries. Therefore, studies that meet just basic criteria for a stroke incidence study set up by Malmgren et al¹⁵ and then updated by Sudlow and Warlow¹⁶ (including studies that accommodate all 3 WHO STEPS surveys and studies based on a special door-to-door survey, as explained above) remain acceptable as basic population-based stroke and TIA incidence studies. There is an obvious lack of even these studies in most parts of the world,^{5,26} and even fewer population-based stroke incidence studies exist on trends in stroke incidence and outcomes.

For stroke incidence studies that use the new definitions of stroke, data on TIA with or without neuroimaging evidence of brain infarct, data on stroke and TIA by age, sex, and ethnic/racial groups should be presented separately and combined. A checklist for standard data reporting is presented in Table 2.

Impact and Implication of New Criteria

As suggested by Sacco et al³¹ changing the definition of stroke and TIA will have significant effects on stroke/TIA surveillance, burden estimates, and prognosis. As the new clinical and tissue-based stroke definition will include formerly diagnosed TIAs with evidence of brain lesion, the number of strokes will be increased, and overall stroke severity reduced. Preliminary estimates suggest that adopting this definition of stroke, for example, in the United States would lower annual incidence rates of TIA by 33%, but increase the rate of ischemic stroke by 33%.^{11,61} It is likely that overall stroke severity will also be reduced by approximately the same amount. However, population-based studies in various populations adopting the new definitions of stroke and TIA are required to estimate the effect of changing stroke and TIA definitions on stroke and TIA incidence and outcomes and to provide new comparable estimates. It is our hope that the wider use of the updated alongside the classical criteria for population-based stroke incidence studies will advance our knowledge on changing stroke epidemiology in the world and further facilitate evidence-based healthcare planning, priority setting, and resource allocation for people with acute cerebrovascular events, and, as a consequence, save millions of lives.

Acknowledgments

Dr Feigin prepared the first draft. Dr Norrving, C.L.M. Sudlow, and Dr Sacco reviewed and edited the draft and approved the final version of the article.

Disclosures

None.

References

1. GBD 2015 Neurological Disorders Collaborator Group. Global, regional, and national burden of neurological disorders during 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Neurol*. 2017;16:877–897. doi: 10.1016/S1474-4422(17)30299-5

2. Feigin VL, Lawes CM, Bennett DA, Barker-Collo SL, Parag V. Worldwide stroke incidence and early case fatality reported in 56 population-based studies: a systematic review. *Lancet Neurol*. 2009;8:355–369. doi: 10.1016/S1474-4422(09)70025-0
3. Feigin VL, Krishnamurthi RV, Parmar P, Norrving B, Mensah GA, Bennett DA, et al: GBD 2013 Writing Group; GBD 2013 Stroke Panel Experts Group. Update on the global burden of ischemic and hemorrhagic stroke in 1990–2013: the GBD 2013 Study. *Neuroepidemiology*. 2015;45:161–176. doi: 10.1159/000441085
4. Global Burden of Disease Study 2013 Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015;386:743–800. doi: 10.1016/S0140-6736(15)60692-4
5. Thrift AG, Thayabaranathan T, Howard G, Howard VJ, Rothwell PM, Feigin VL, et al. Global stroke statistics. *Int J Stroke*. 2017;12:13–32. doi: 10.1177/1747493016676285
6. Gutierrez J, Williams OA. A decade of racial and ethnic stroke disparities in the United States. *Neurology*. 2014;82:1080–1082. doi: 10.1212/WNL.0000000000000237
7. Wang Y, Rudd AG, Wolfe CD. Age and ethnic disparities in incidence of stroke over time: the South London Stroke Register. *Stroke*. 2013;44:3298–3304. doi: 10.1161/STROKEAHA.113.002604
8. Sacco RL, Gardener H, Wang K, Dong C, Ciliberti-Vargas MA, Gutierrez CM, et al. Racial-ethnic disparities in acute stroke care in the Florida-Puerto Rico collaboration to reduce stroke disparities study. *J Am Heart Assoc*. 2017;6:e004073. doi: 10.1161/JAHA.116.004073
9. GBD 2016 Risk Factors Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017;390:1345–1422. doi: 10.1016/S0140-6736(17)32366-8
10. Feigin VL, Krishnamurthi RV, Barker-Collo S, McPherson KM, Barber PA, Parag V, et al: ARCOS IV Group. 30-year trends in stroke rates and outcome in Auckland, New Zealand (1981–2012): a multi-ethnic population-based series of studies. *PLoS One*. 2015;10:e0134609. doi: 10.1371/journal.pone.0134609
11. Ovbiagele B, Nguyen-Huynh MN. Stroke epidemiology: advancing our understanding of disease mechanism and therapy. *Neurotherapeutics*. 2011;8:319–329. doi: 10.1007/s13311-011-0053-1
12. Bogiatzi C, Hackam DG, McLeod AI, Spence JD. Secular trends in ischemic stroke subtypes and stroke risk factors. *Stroke*. 2014;45:3208–3213. doi: 10.1161/STROKEAHA.114.006536
13. McCarron MO, Davey Smith G, McCarron P. Secular stroke trends: early life factors and future prospects. *QJM*. 2006;99:117–122. doi: 10.1093/qjmed/hcl008
14. Aho K, Harmsen P, Hatano S, Marquardsen J, Smirnov VE, Strasser T. Cerebrovascular disease in the community: results of a WHO collaborative study. *Bull World Health Organ*. 1980;58:113–130.
15. Malmgren R, Warlow C, Bamford J, Sandercock P. Geographical and secular trends in stroke incidence. *Lancet*. 1987;2:1196–1200.
16. Sudlow CL, Warlow CP. Comparing stroke incidence worldwide: what makes studies comparable? *Stroke*. 1996;27:550–558.
17. Feigin VL, Carter K. Editorial comment—stroke incidence studies one step closer to the elusive gold standard? *Stroke*. 2004;35:2045–2047.
18. Feigin V, Hoorn SV. How to study stroke incidence. *Lancet*. 2004;363:1920. doi: 10.1016/S0140-6736(04)16436-2
19. Truelsen T, Bonita R, Jamrozik K. Surveillance of stroke: a global perspective. *Int J Epidemiol*. 2001;30(suppl 1):S11–S16.
20. Feigin VL. Stroke in developing countries: can the epidemic be stopped and outcomes improved? *Lancet Neurol*. 2007;6:94–97. doi: 10.1016/S1474-4422(07)70007-8
21. Easton JD, Saver JL, Albers GW, Alberts MJ, Chaturvedi S, Feldmann E, et al; American Heart Association; American Stroke Association Stroke Council; Council on Cardiovascular Surgery and Anesthesia; Council on Cardiovascular Radiology and Intervention; Council on Cardiovascular Nursing; Interdisciplinary Council on Peripheral Vascular Disease. Definition and evaluation of transient ischemic attack: a scientific statement for healthcare professionals from the American Heart Association/American Stroke Association Stroke Council; Council on Cardiovascular Surgery and Anesthesia; Council on Cardiovascular Radiology and Intervention; Council on Cardiovascular Nursing; and the Interdisciplinary Council on Peripheral Vascular Disease. The American Academy of Neurology affirms the value of this statement as

- an educational tool for neurologists. *Stroke*. 2009;40:2276–2293. doi: 10.1161/STROKEAHA.108.192218
22. World Health Organization. ICD-11 Beta Draft (Mortality and Morbidity Statistics). <https://icd.who.int/dev11l-m/en>. Accessed March 26, 2018.
 23. Longstreth WT Jr, Gasca NC, Gottesman RF, Pearce JB, Sacco RL. Adjudication of transient ischemic attack and stroke in the multi-ethnic study of atherosclerosis. *Neuroepidemiology*. 2018;50:23–28. doi: 10.1159/000486174
 24. Nadarajan V, Perry RJ, Johnson J, Werring DJ. Transient ischaemic attacks: mimics and chameleons. *Pract Neurol*. 2014;14:23–31. doi: 10.1136/practneurol-2013-000782
 25. Rothwell PM, Coull AJ, Silver LE, Fairhead JF, Giles MF, Lovelock CE, et al; Oxford Vascular Study. Population-based study of event-rate, incidence, case fatality, and mortality for all acute vascular events in all arterial territories (Oxford Vascular Study). *Lancet*. 2005;366:1773–1783. doi: 10.1016/S0140-6736(05)67702-1
 26. Barber PA, Krishnamurthi R, Parag V, Anderson NE, Ranta A, Kilfoyle D, et al; ARCOS IV Study Group. Incidence of transient ischemic attack in Auckland, New Zealand, in 2011 to 2012. *Stroke*. 2016;47:2183–2188. doi: 10.1161/STROKEAHA.116.014010
 27. Bahit MC, Coppola ML, Riccio PM, Cipriano LE, Roth GA, Lopes RD, et al. First-ever stroke and transient ischemic attack incidence and 30-day case-fatality rates in a population-based study in Argentina. *Stroke*. 2016;47:1640–1642. doi: 10.1161/STROKEAHA.116.013637
 28. Fonseca PG, Weiss PA, Harger R, Moro CH, Longo AL, Gonçalves AR, et al. Transient ischemic attack incidence in Joinville, Brazil, 2010: a population-based study. *Stroke*. 2012;43:1159–1162. doi: 10.1161/STROKEAHA.111.645101
 29. Cancelli I, Janes F, Gigli GL, Perelli A, Zanchettin B, Canal G, et al. Incidence of transient ischemic attack and early stroke risk: validation of the ABCD2 score in an Italian population-based study. *Stroke*. 2011;42:2751–2757. doi: 10.1161/STROKEAHA.110.612705
 30. Béjot Y, Brenière C, Graber M, Garnier L, Durier J, Blanc-Labarre C, et al. Contemporary epidemiology of transient ischemic attack in Dijon, France (2013–2015). *Neuroepidemiology*. 2017;49:135–141. doi: 10.1159/000484638
 31. Sacco RL, Kasner SE, Broderick JP, Caplan LR, Connors JJ, Culebras A, et al; American Heart Association Stroke Council; Council on Cardiovascular Surgery and Anesthesia; Council on Cardiovascular Radiology and Intervention; Council on Cardiovascular and Stroke Nursing; Council on Epidemiology and Prevention; Council on Peripheral Vascular Disease; Council on Nutrition, Physical Activity and Metabolism. An updated definition of stroke for the 21st century: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2013;44:2064–2089. doi: 10.1161/STR.0b013e318296aeca
 32. Adams HP Jr, Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke*. 1993;24:35–41.
 33. Hart RG, Diener HC, Coutts SB, Easton JD, Granger CB, O'Donnell MJ, et al; Cryptogenic Stroke/ESUS International Working Group. Embolic strokes of undetermined source: the case for a new clinical construct. *Lancet Neurol*. 2014;13:429–438. doi: 10.1016/S1474-4422(13)70310-7
 34. Hart RG, Catanese L, Perera KS, Ntaios G, Connolly SJ. Embolic stroke of undetermined source: a systematic review and clinical update. *Stroke*. 2017;48:867–872. doi: 10.1161/STROKEAHA.116.016414
 35. Ay H, Benner T, Arsava EM, Furie KL, Singhal AB, Jensen MB, et al. A computerized algorithm for etiologic classification of ischemic stroke: the Causative Classification of Stroke System. *Stroke*. 2007;38:2979–2984. doi: 10.1161/STROKEAHA.107.490896
 36. McArdle PF, Kittner SJ, Ay H, Brown RD Jr, Meschia JF, Rundek T, et al; NINDS SIGN Study. Agreement between TOAST and CCS ischemic stroke classification: the NINDS SIGN study. *Neurology*. 2014;83:1653–1660. doi: 10.1212/WNL.0000000000000942
 37. Lindley RI, Warlow CP, Wardlaw JM, Dennis MS, Slattery J, Sandercock PA. Interobserver reliability of a clinical classification of acute cerebral infarction. *Stroke*. 1993;24:1801–1804.
 38. National Institute of Health, National Institute of Neurological Disorders and Stroke. Stroke Scale. https://www.ninds.nih.gov/sites/default/files/NIH_Stroke_Scale_Booklet.pdf. Accessed May 3, 2018.
 39. Williams LS, Yilmaz EY, Lopez-Yunez AM. Retrospective assessment of initial stroke severity with the NIH Stroke Scale. *Stroke*. 2000;31:858–862.
 40. Krishnamurthi RV, Barker-Collo S, Parag V, Parmar P, Witt E, Jones A, et al. Stroke incidence by major pathological type and ischemic subtypes in the Auckland regional community stroke studies: changes between 2002 and 2011. *Stroke*. 2018;49:3–10. doi: 10.1161/STROKEAHA.117.019358
 41. Hajat C, Heuschmann PU, Coshall C, Padayachee S, Chambers J, Rudd AG, et al. Incidence of aetiological subtypes of stroke in a multi-ethnic population based study: the South London Stroke Register. *J Neurol Neurosurg Psychiatry*. 2011;82:527–533. doi: 10.1136/jnnp.2010.222919
 42. Schulz UG, Rothwell PM. Differences in vascular risk factors between etiological subtypes of ischemic stroke: importance of population-based studies. *Stroke*. 2003;34:2050–2059. doi: 10.1161/01.STR.0000079818.08343.8C
 43. Petty GW, Brown RD Jr, Whisnand JP, Sicks JD, O'Fallon WM, Wiebers DO. Ischemic stroke subtypes: a population-based study of incidence and risk factors. *Stroke*. 1999;30:2513–2516.
 44. Kolominsky-Rabas PL, Weber M, Gefeller O, Neundoerfer B, Heuschmann PU. Epidemiology of ischemic stroke subtypes according to TOAST criteria: incidence, recurrence, and long-term survival in ischemic stroke subtypes: a population-based study. *Stroke*. 2001;32:2735–2740.
 45. Palm F, Urbanek C, Wolf J, Bugge F, Kleemann T, Hennerici MG, et al. Etiology, risk factors and sex differences in ischemic stroke in the Ludwigshafen Stroke Study, a population-based stroke registry. *Cerebrovasc Dis*. 2012;33:69–75. doi: 10.1159/000333417
 46. Corso G, Bottacchi E, Giardini G, Di Giovanni M, Meloni T, Pesenti Campagnoni M, et al. Epidemiology of stroke in northern Italy: the Cerebrovascular Aosta Registry, 2004–2008. *Neurol Sci*. 2013;34:1071–1081. doi: 10.1007/s10072-012-1185-8
 47. Díaz-Guzmán J, Egido JA, Gabriel-Sánchez R, Barberá-Comes G, Fuentes-Gimeno B, Fernández-Pérez C; IBERICTUS Study Investigators of the Stroke Project of the Spanish Cerebrovascular Diseases Study Group. Stroke and transient ischemic attack incidence rate in Spain: the IBERICTUS study. *Cerebrovasc Dis*. 2012;34:272–281. doi: 10.1159/000342059
 48. Leyden JM, Kleinig TJ, Newbury J, Castle S, Cranfield J, Anderson CS, et al. Adelaide stroke incidence study: declining stroke rates but many preventable cardioembolic strokes. *Stroke*. 2013;44:1226–1231. doi: 10.1161/STROKEAHA.113.675140
 49. White H, Boden-Albala B, Wang C, Elkind MS, Rundek T, Wright CB, et al. Ischemic stroke subtype incidence among whites, blacks, and Hispanics: the Northern Manhattan Study. *Circulation*. 2005;111:1327–1331. doi: 10.1161/01.CIR.0000157736.19739.D0
 50. Taub NA, Lemic-Stojcevic N, Wolfe CD. Capture-recapture methods for precise measurement of the incidence and prevalence of stroke. *J Neurol Neurosurg Psychiatry*. 1996;60:696–697.
 51. Tilling K, Sterne JA, Wolfe CD. Estimation of the incidence of stroke using a capture-recapture model including covariates. *Int J Epidemiol*. 2001;30:1351–1359; discussion 1359.
 52. Ganesh A, Luengo-Fernandez R, Wharton RM, Gutnikov SA, Silver LE, Mehta Z, et al. Time course of evolution of disability and cause-specific mortality after ischemic stroke: implications for trial design. *J Am Heart Assoc*. 2017;6:e005788. doi: 10.1161/JAHA.117.005788
 53. Eriksson M, Norrving B, Terént A, Stegmayr B. Functional outcome 3 months after stroke predicts long-term survival. *Cerebrovasc Dis*. 2008;25:423–429. doi: 10.1159/000121343
 54. Rothwell PM, Coull AJ, Giles MF, Howard SC, Silver LE, Bull LM, et al; Oxford Vascular Study. Change in stroke incidence, mortality, case-fatality, severity, and risk factors in Oxfordshire, UK from 1981 to 2004 (Oxford Vascular Study). *Lancet*. 2004;363:1925–1933. doi: 10.1016/S0140-6736(04)16405-2
 55. Lindsay P, Furie KL, Davis SM, Donnan GA, Norrving B. World Stroke Organization global stroke services guidelines and action plan. *Int J Stroke*. 2014;9(suppl A100):4–13. doi: 10.1111/ijss.12371
 56. Rocca WA, Reggioni A, Savettieri G, Salemi G, Patti F, Meneghini F, et al. Stroke incidence and survival in three Sicilian municipalities. Sicilian Neuro-Epidemiologic Study (SNES) Group. *Ital J Neurol Sci*. 1998;19:351–356.
 57. Li SC, Schoenberg BS, Wang CC, Cheng XM, Bolis CL, Wang KJ. Cerebrovascular disease in the People's Republic of China: epidemiologic and clinical features. *Neurology*. 1985;35:1708–1713.
 58. Wang W, Jiang B, Sun H, Ru X, Sun D, Wang L, et al; NESS-China Investigators. Prevalence, incidence, and mortality of stroke in China: results from a nationwide population-based survey of 480687 adults.

- Circulation*. 2017;135:759–771. doi: 10.1161/CIRCULATIONAHA.116.025250
59. Bennett DA, Brayne C, Feigin VL, Barker-Collo S, Brainin M, Davis D, et al. Explanation and elaboration of the standards of reporting of neurological disorders checklist: a guideline for the reporting of incidence and prevalence studies in neuroepidemiology. *Neuroepidemiology*. 2015;45:113–137. doi: 10.1159/000439132
60. Bennett DA, Brayne C, Feigin VL, Barker-Collo S, Brainin M, Davis D, et al. Development of the standards of reporting of neurological disorders (STROND) checklist: a guideline for the reporting of incidence and prevalence studies in neuroepidemiology. *Eur J Epidemiol*. 2015;30:569–576. doi: 10.1007/s10654-015-0034-5
61. B ejot Y, Mehta Z, Giroud M, Rothwell PM. Impact of completeness of ascertainment of minor stroke on stroke incidence: implications for ideal study methods. *Stroke*. 2013;44:1796–1802. doi: 10.1161/STROKEAHA.113.000949

KEY WORDS: cause of death ■ epidemiology ■ risk factors ■ stroke ■ subarachnoid hemorrhage



Stroke